

Activation of Non-Local Quantum Resources

Miguel Navascués¹ and Tamás Vértesi²

¹*Facultad de Matemáticas, Universidad Complutense de Madrid*

²*Institute of Nuclear Research of the Hungarian Academy of Sciences*

We find two two-qubit bipartite states ρ_1, ρ_2 such that arbitrarily many copies of one or the other cannot exhibit non-local correlations in a two settings/two outcomes Bell scenario. However, the bipartite state $\rho_1 \otimes \rho_2$ violates the Clauser-Horne-Shimony-Holt (CHSH) Bell inequality [12] by an amount of 2.023. We also identify a CHSH-local state ρ such that $\rho^{\otimes 2}$ is CHSH-violating. The tools employed can be easily adapted to find instances of non-locality activation in arbitrary Bell scenarios.

PACS numbers:

In 1964 John S. Bell refuted the Einstein-Podolski-Rosen (EPR) argument against Quantum Mechanics [1] by showing that certain correlations generated by distant observers measuring a quantum system are impossible to reproduce in any local realistic theory [2]. Such correlations, which do not arise as marginals of a greater probability distribution, have been termed ‘non-local’ in recent years. Initially a foundational concept, non-locality has lately found applications in several areas of Quantum Information, like quantum cryptography [3], certified randomness generation [4] and device-independent quantum state estimation [5, 6]. In any concrete implementation of the former protocols, non-locality is quantified according to some linear figure of merit, the so-called Bell functional. The amount by which the Bell value of a given quantum set of correlations differs from the classical maximum (the Bell inequality violation) shall thus be regarded as a resource in these primitives.

A prerequisite for the emergence of non-locality is the presence of entanglement in the quantum state where the measurements are carried out. However, the opposite implication is far from clear. Although there exist examples of entangled quantum systems unable to produce non-local correlations [7], Popescu and Gisin showed that some of those states exhibit a non-local behavior if, prior to the Bell tests, they are subject to local measurements and post-selected according to the measurement results [8, 9]. Some time later, Peres observed that an even bigger set of local states reveal a non-local nature when post-selections and measurements are conducted, not over a single local state, but over several identical copies [10].

In view of Peres’ results, it is reasonable to ask whether this form of ‘non-locality activation’ could also be achieved by performing measurements over ensembles of local states only, without resorting to any form of post-selection. This corresponds to a natural situation where the distant parties receive a number of local states and are asked to violate a Bell inequality with no prior communication. Although there have been some numerical attempts to attack this problem [11], it is still an open question whether such a ‘tensoring-mediated activation’ exists.

If our figure of merit is not mere deviation from lo-

cality, but the violation of a specific Bell inequality, the corresponding scenario is quite similar to the one above: given a Bell functional $\mathcal{B}$, we can define $\mathcal{B}$ -local states as those quantum systems where the evaluation of $\mathcal{B}$ cannot exceed the classical bound. Analogously, given a set of $\mathcal{B}$ -local states, we may speak of $\mathcal{B}$ -activation if those states can be locally engineered to produce a $\mathcal{B}$ violation.

When $\mathcal{B}$ happens to be the Clauser-Horne-Shimony-Holt (CHSH) inequality [12] and local filtering is allowed, the connection between entanglement and non-locality is well understood. We know that the set of quantum states which violate CHSH under local post-selection of many copies coincides with the set of distillable states [13]. Moreover, we know that for any entangled state σ_1 there exists another state σ_2 none of whose local filterings violates the CHSH inequality, but such that $\sigma_1 \otimes \sigma_2$ violates CHSH after some local post-selection [14]. Choosing σ_1 undistillable (and so unable to violate CHSH) in the above proposition, we thus have that CHSH non-locality can be activated by post-selecting tensor products of CHSH-local states.

Nothing is known, though, about activation of bipartite Bell inequalities by tensoring alone (note, however, the recent work [15], where the authors show instances of non-locality activation in the multipartite setting). This led Liang to ask in [11, 16] whether there exist two CHSH-local states ρ_1, ρ_2 such that $\rho_1 \otimes \rho_2$ generates CHSH-violating statistics when measured in the appropriate bases.

In this article we will answer this question in two different ways: first, we will show the existence of two states ρ_1, ρ_2 such that $\rho_1^{\otimes N}$ and $\rho_2^{\otimes N}$ do not violate CHSH for any N , but $\rho_1 \otimes \rho_2$ does. Second, we will construct a CHSH-local state ρ such that $\rho^{\otimes 2}$ is CHSH-violating.

Some examples of non-locality activation

Consider a situation where two space-like separated observers, say, Alice and Bob, are conducting measurements on a joint quantum system. Whenever Alice applies an interaction $x \in \{1, \dots, s_A\}$ on her subsystem, she will read an outcome $a \in \{1, \dots, d_A\}$ in her detector; likewise, if Bob interacts with his subsystem in some way $y \in \{1, \dots, s_B\}$, his measurement devices will register a value $b \in \{1, \dots, d_B\}$. Varying the choice of measurement

settings x, y , and after many repetitions of the experiment, Alice and Bob would be able to estimate the set of correlations $\{P(a, b|x, y)\}$, where $P(a, b|x, y)$ denotes the probability that Alice and Bob observe the results a and b when they respectively perform the interactions x and y . Given such a $s_{ASB}d_{ADB}$ Bell scenario, we will say that $\{P(a, b|x, y)\}$ is local iff there exists a global measure $\mathcal{P}(a_1, \dots, a_{s_A}, b_1, \dots, b_{s_B})$ such that

$$P(a, b|x, y) = \mathcal{P}(a_x = a, b_y = b). \quad (1)$$

The 2222 Bell scenario is the simplest bipartite setting where one can expect to find locality violations. There, the locality of a bipartite distribution $\{P(a, b|x, y) : x, y, a, b = 1, 2\}$ can be decided by checking the value of the Clauser-Horne-Shimony-Holt (CHSH) inequality [12]. More concretely, modulo permutations of the settings, $P(a, b|x, y)$ is local iff

$$|\langle M^1 N^1 \rangle + \langle M^1 N^2 \rangle + \langle M^2 N^1 \rangle - \langle M^2 N^2 \rangle| \leq 2, \quad (2)$$

where M^x (N^y) is any observable that assigns values $+1$ or -1 to the two possible outcomes of Alice's (Bob's) measurement x (y). Consequently, any bipartite state ρ is local in the 2222 scenario iff, for any set of operators $\{M^1, M^2, N^1, N^2\}$ with spectrum $\{-1, 1\}$, the Bell operator

$$\mathcal{B} \equiv M^1 \otimes N^1 + M^2 \otimes N^1 + M^1 \otimes N^2 - M^2 \otimes N^2 \quad (3)$$

is such that $|\text{tr}(\mathcal{B}\rho)| \leq 2$ [23].

The bipartite quantum states ρ_1, ρ_2 alluded to in the abstract of the paper, as well as the rest of the states and measurements referred to along this article, can be found in [17]. It can be checked that $\rho_1^{AB} = \text{tr}_{B'}(\rho_1^{ABB'})$, $\rho_2^{AB} = \text{tr}_{A'}(\rho_2^{AA'B})$, for some tripartite states $\rho_1^{ABB'}$ and $\rho_2^{AA'B}$. These states in turn satisfy $\rho_1^{ABB'} = \rho_1^{A'B'B}$ and $\rho_2^{AA'B} = \rho_2^{A'AB}$, i.e., $\rho_1^{ABB'}$ and $\rho_2^{AA'B}$ are permutationally invariant with respect to the systems B, B' and A, A' , respectively. Using modern terminology, we would say that ρ_1 (ρ_2) admits a 2-symmetric extension with respect to system B (A) [18]. In the following, we will denote by S_C^2 the set of all quantum states admitting a 2-extension with respect to system C .

As observed in [19], any state $\sigma^{AB} \in B(H \otimes H)$ that admits a 2 symmetric extension (say, with respect to system B) cannot violate any Bell inequality involving 2 measurement settings on Bob's side. Indeed, denote Bob's measurements by $\{y_1, y_2\}$, and let $\{E_x^a, F_{b_1}^{y_1}, F_{b_2}^{y_2}\}$ be complete systems of Positive Operator Valued Measure (POVM) elements, with $x = 1, \dots, s$. Then, one can check that the set of bipartite correlations $p(a, b|x, y) \equiv \text{tr}(E_x^a \otimes F_b^y \sigma^{AB})$ derives from the global probability distribution

$$P(a_1, a_2, \dots, a_s, b_1, b_2) \equiv p(b_1, b_2) \prod_{x=1}^s p(a_x|x, b_1, b_2), \quad (4)$$




FIG. 1: Left: 2-extendable states with respect to a given system (say, 2) are invariant under tensor products; this is so because the tensor product of their extensions is invariant under the interchange of the systems 2 and 3. Right: tensor product of two states admitting 2 symmetric extensions to different systems. In general, these composite states do not admit a 2 symmetric extension.

where $p(b_1, b_2) = \text{tr}(\sigma^{BB'} F_{b_1}^{y_1} \otimes F_{b_2}^{y_2})$ and

$$P(a|x, b_1, b_2) \equiv \frac{\text{tr}(\sigma^{ABB'} E_a^x \otimes F_{b_1}^{y_1} \otimes F_{b_2}^{y_2})}{p(b_1, b_2)}. \quad (5)$$

As sketched in Figure 1 (left), if we tensor two states $\rho_1, \rho_2 \in S_B^2$ (S_A^2), the resulting state $\rho_1 \otimes \rho_2$ can also be extended symmetrically with respect to system B (A). It follows that arbitrarily many copies of ρ_1 (ρ_2) can never violate any Bell inequality involving two settings on Bob's (Alice's) side.

What happens when we join ρ_1 and ρ_2 ? Following Figure 1, if $\rho_1 \in S_B^2$ and $\rho_2 \in S_A^2$, then the state $\rho_1 \otimes \rho_2$ does not necessarily admit a 2-symmetric extension, and so (in principle) could violate CHSH. And, actually, one can check that the measurement operators given in [17], applied to $\rho_1^{AB} \otimes \rho_2^{A'B'}$, result in a CHSH violation of 2.02324.

In order to come up with the previous example, we used the fact that, for fixed (M^x, N^y, ρ_1) , finding a 2-extendable state ρ_2 that maximizes the CHSH value of the system $(M^x, N^y, \rho_1, \rho_2)$ is a trivial exercise of linear algebra. In effect, define $\mathcal{B} = M_{AA'}^1 \otimes N_{BB'}^1 + M_{AA'}^1 \otimes N_{BB'}^2 + M_{AA'}^2 \otimes N_{BB'}^1 - M_{AA'}^2 \otimes N_{BB'}^2$ and $\mathcal{B}_2 \equiv \text{tr}_{AB}(\rho_1^{AB} \otimes \mathbb{I}_{A'B'} \mathcal{B})$, and let V_{GH} denote the operator that swaps systems G and H . Then,

$$\begin{aligned} \max\{\text{tr}(\mathcal{B}\rho_1 \otimes \rho_2) : \rho_2^{A'B'} \in S_{A'}^2\} &= \\ = \max\{\text{tr}(\mathcal{B}_2\rho_2) : \rho_2^{A'B'} \in S_{A'}^2\} &= \\ = \max\{\text{tr}(\tilde{\mathcal{B}}_2\rho^{CA'B'}) : \rho^{CA'B'}\}, \end{aligned} \quad (6)$$

where $\rho^{CA'B'}$ is an arbitrary normalized quantum state and $\tilde{\mathcal{B}}_2 \in B(H_C \otimes H_{A'} \otimes H_{B'})$ is given by

$$\tilde{\mathcal{B}}_2 \equiv \frac{1}{2} \{ \mathbb{I}_C \otimes \mathcal{B}_2 + V_{CA'} (\mathbb{I}_C \otimes \mathcal{B}_2) V_{CA'} \}. \quad (7)$$

The problem just reduces to finding the greatest eigenvalue of $\tilde{\mathcal{B}}_2$. The corresponding eigenvector $|\psi\rangle$ can then be transformed into a 2-extendable state ρ_2 through $\rho_2 = \text{tr}_C(V_{CA'}|\psi\rangle\langle\psi|V_{CA'} + |\psi\rangle\langle\psi|)/2$. Likewise, for fixed M^x, M^y, ρ_2 , the optimal ρ_1 can be found efficiently.

Conversely, if (M^x, ρ_1, ρ_2) are fixed, one can find the observables on Bob's side that maximize the CHSH value using the procedure sketched in [20]. That is, given

$$\begin{aligned} F^1 &\equiv \text{tr}_{AA'} \{ ([M^1 + M^2] \otimes \mathbb{I}_{BB'}) (\rho_1 \otimes \rho_2) \}, \\ F^2 &\equiv \text{tr}_{AA'} \{ ([M^1 - M^2] \otimes \mathbb{I}_{BB'}) (\rho_1 \otimes \rho_2) \}, \end{aligned} \quad (8)$$

the problem consists in maximizing $\text{tr}\{N^1 F^1 + N^2 F^2\}$ over all operators $-\mathbb{I}_{BB'} \leq N^k \leq \mathbb{I}_{BB'}, k = 1, 2$. Now, let $\sum_j \lambda_j^k |\phi_j^k\rangle\langle\phi_j^k|$ be the singular value decomposition of F^k . Then, the optimal N^1, N^2 are $N^k = \sum_j \text{sgn}(\lambda_j^k) |\phi_j^k\rangle\langle\phi_j^k|$. Analogously, we can easily obtain the optimal M^1, M^2 for fixed N^1, N^2, ρ_1, ρ_2 .

Starting with random states and operators, one can then optimize one set of variables at a time, following a cycle like $(M^1, M^2) \rightarrow \rho_1 \rightarrow (N^1, N^2) \rightarrow \rho_2 \rightarrow (M^1, M^2) \rightarrow \dots$. This optimization procedure frequently gets stuck in local maxima, so it has to be repeated several times before reaching global optimality.

We repeated the whole scheme with pairs of states in dimensions $d = 3, 4, 5$, but we were not able to overcome the activation value 2.040167, which is attained by two pairs of qutrits.

Notice that the former tools can be used to find instances of activation of any two-setting Bell inequality. If the number of outcomes of such inequality is greater than two, though, it becomes necessary to use semidefinite programming (SDP) [21] in order to perform the corresponding POVM optimizations. The resulting programs are therefore much slower, and wide searches over high dimensions become computationally expensive. Nevertheless, we managed to find two states of dimension $3 \otimes 3$ that, although unable to violate the three-outcome CGLMP inequality [22] separately, could give rise to a CGLMP value of 2.030126 when taken together.

Symmetries

All states found by our method are highly asymmetric by construction. Suppose, on the contrary, that we are interested in finding symmetric states ρ_1, ρ_2 with the property that ρ_1, ρ_2 do not violate two-setting Bell inequalities, but such that $\rho_1 \otimes \rho_2$ violates a certain two-setting Bell inequality. Then one could try the scheme presented in Figure 2, i.e., both ρ_1 and ρ_2 are the result of symmetrizing bipartite states $\rho_1^{AB}, \rho_2^{AB} \in S_B^2$. Being a mixture of $2n$ -local and $n2$ -local states, such states are 22 -local.

Like before, it is not difficult to reduce the problem of maximizing $\text{tr}(M\rho)$ over all possible such states to a




FIG. 2: Two bipartite states resulting from the symmetrization of states in S_B^2 (or, equivalently, S_A^2) are not necessarily local in two-setting scenarios due to the presence of mixed terms admitting different types of extensions.

diagonalization problem. Using this new procedure, we were not able to find instances of symmetric activation in dimensions 2 and 3. However, we managed to find two 22-local symmetric states $\rho'_1, \rho'_2 \in B(\mathbb{C}^4 \otimes \mathbb{C}^4)$ with the property that $\rho'_1 \otimes \rho'_2$ violates CHSH by 2.01159. Note, though, that this time locality does not extend to multiple copies of the states. That is, in principle $\rho_1^{\otimes N}$ could be CHSH-violating for some N .

This new bound, however, can be beaten easily: take the previous two-qubit asymmetric states ρ_1, ρ_2 that produce a CHSH violation of 2.02324 when taken together, and let (M^x, N^y) be the necessary measurements to ensure so. If, in addition to these systems, Alice and Bob have access to two extra ancillary qubits, then we can take the new symmetric states to be

$$\tilde{\rho}_{1,2} = \frac{1}{2} (|0\rangle\langle 0|_A \otimes |1\rangle\langle 1|_B \otimes \rho_{1,2} + |1\rangle\langle 1|_A \otimes |0\rangle\langle 0|_B \otimes V_{AB} \rho_{1,2} V_{AB}). \quad (9)$$

Clearly, these states cannot violate the CHSH inequality. On the other hand, it is easy to check that Alice and Bob can violate CHSH if each of them measures its part of the state $\tilde{\rho}_1 \otimes \tilde{\rho}_2$ according to the following measurement scheme: let $z \in \{1, 2\}$ be the label of the measurement to be implemented. First, measure the ancillary qubits in the computational basis. If both qubits are in state $|0\rangle$, measure M^z in the rest of the system and output the result (+1 or -1); if the ancillary qubits are both in state $|1\rangle$, measure N^z and output the result; if they are in different states, output 1.

With this procedure, Alice and Bob should observe a CHSH violation of $(2 \times 2.02324 + 2 \times 2)/4 = 2.01162$.

More generally, from any pair of CHSH-local states $\sigma_1, \sigma_2 \in B(\mathbb{C}^d \otimes \mathbb{C}^d)$ such that $\sigma_1 \otimes \sigma_2$ violates CHSH by an amount $2 + \Delta$, one can derive two CHSH-local symmetric states $\tilde{\sigma}_1, \tilde{\sigma}_2 \in B(\mathbb{C}^{2d} \otimes \mathbb{C}^{2d})$ violating CHSH by $2 + \Delta/2$.

This scheme of using simple instances of non-locality activation to construct more sophisticated ones can be carried further. Suppose that we are given any such pair of states σ_1, σ_2 . Then one can also find a state

$\tilde{\sigma} \in B(\mathbb{C}^{2d} \otimes \mathbb{C}^{2d})$ such that $\tilde{\sigma}$ does not violate CHSH, but $\tilde{\sigma}^{\otimes 2}$ exhibits a CHSH violation of $2 + \Delta/2$. Indeed, take the state to be

$$\tilde{\sigma} \equiv \frac{1}{2}(|1\rangle\langle 1|_A \otimes |1\rangle\langle 1|_B \otimes \sigma_1 + |0\rangle\langle 0|_A \otimes |0\rangle\langle 0|_B \otimes \sigma_2), \quad (10)$$

and, as before, let Alice (Bob) measure her (his) main subsystem with M^x , $V_{AA'}M^xV_{AA'}$ or $\mathbb{I}_{AA'}$ (N^y , $V_{BB'}N^yV_{BB'}$ or $\mathbb{I}_{BB'}$) depending on the state of her (his) ancillary qubits.

Combining both constructions, it is clear that, starting from any CHSH-activating pair $\sigma_1, \sigma_2 \in B(\mathbb{C}^d \otimes \mathbb{C}^d)$, we can build a CHSH-local symmetric state $\hat{\sigma} \in B(\mathbb{C}^{4d} \otimes \mathbb{C}^{4d})$ such that $\hat{\sigma}^{\otimes 2}$ violates CHSH by an amount $2 + \Delta/4$.

The above results can be easily generalized to arbitrary symmetric Bell inequalities.

From the previous section, it thus follows that there exists a 22-local symmetric state $\rho \in B(\mathbb{C}^8 \otimes \mathbb{C}^8)$ ($\rho' \in B(\mathbb{C}^{12} \otimes \mathbb{C}^{12})$) with the property that two copies of it can violate the CHSH inequality by an amount of 2.00581 (2.01). Likewise, there exists a CGLMP-local symmetric state $\rho'' \in B(\mathbb{C}^{12} \otimes \mathbb{C}^{12})$ such that $(\rho'')^{\otimes 2}$ violates the three-outcome CGLMP inequality by 2.007531.

Conclusion

In this communication we have given a systematic procedure to construct examples of non-locality activation. By exploiting the concept of 2-extendibility, we were able

to find two two-qubit bipartite states ρ_1, ρ_2 such that even though neither $\rho_1^{\otimes N}$ nor $\rho_2^{\otimes N}$ violates CHSH, $\rho_1 \otimes \rho_2$ produces a CHSH parameter of 2.023. We have thus solved problem #21 of the Hannover List of Open Problems in Quantum Information [16].

We have explained how the previous results can be used to find examples of CHSH self-activation, i.e., CHSH-local states $\tilde{\sigma}$ such that $\tilde{\sigma}^{\otimes 2}$ is CHSH-violating. Finally, we have also shown how to extend our approach to other 2-setting Bell inequalities, like the CGLMP family. Clearly, there is no need to stop here: our techniques can be easily adapted to find examples of activation of non-locality in scenarios with more than two settings per site and more than two sites.

It would be interesting to find out if the small deviations of locality found here are close to optimal, or, on the contrary, non-locality activation can be as huge as we want provided that we go to dimensions high enough. Perhaps the fact that the maximum activation values for CHSH and CGLMP seem to be the same could give us some clue here. We also wonder whether an extreme extension of our results could be given, i.e., whether one could find states ρ_1, ρ_2 such that $\rho_1 \otimes \rho_2$ is non-local, but neither $\rho_1^{\otimes N}$ nor $\rho_2^{\otimes N}$ violate *any* Bell inequality.

Acknowledgements

M. N. has been supported by the European project QUEVADIS. T.V. has been supported by a János Bolyai Grant of the Hungarian Academy of Sciences.

-
- [1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] J. S. Bell, Physics **1**, 195 (1964).
 - [3] A. Acín, N. Brunner, N. Gisin, S. Massar, S. Pironio and V. Scarani, Phys. Rev. Lett. **98**, 230501 (2007).
 - [4] S. Pironio, A. Acín, S. Massar, A. Boyer de la Giroday, D. N. Matsukevich, P. Maunz, S. Olmschenk, D. Hayes, L. Luo, T. A. Manning and C. Monroe, Nature **464**, 1021 (2010).
 - [5] C.-E. Bardyn, T. C. H. Liew, S. Massar, M. McKague and V. Scarani, Phys. Rev. A **80**, 062327 (2009).
 - [6] M. McKague, and M. Mosca, arXiv:1006.0150v1.
 - [7] R. F. Werner, Phys. Rev. A **40**, 4277 (1989); J. Barrett, Phys. Rev. A **65**, 042302 (2002); A. Acín, N. Gisin, and B. Toner, Phys. Rev. A **73**, 062105 (2006); M. L. Almeida, S. Pironio, J. Barrett, G. Toth, A. Acín, Phys. Rev. Lett. **99**, 040403 (2007).
 - [8] S. Popescu, Phys. Rev. Lett. **74**, 2619 (1995).
 - [9] N. Gisin, Phys. Lett. A **210**, 151 (1996).
 - [10] A. Peres, Phys. Rev. A **54**, 2685 (1996).
 - [11] Y.-C. Liang and A. C. Doherty, Phys. Rev. A, **73**, 052116 (2006).
 - [12] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
 - [13] Ll. Masanes, Phys. Rev. Lett. **97**, 050503 (2006).
 - [14] Ll. Masanes, Y.-C. Liang and A. C. Doherty, Phys. Rev. Lett. **100**, 090403 (2008).
 - [15] D. Cavalcanti, M. Almeida, A. Acín and V. Scarani, arXiv:1010.0900.
 - [16] O. Krueger and R. F. Werner, arXiv:quant-ph/0504166.
 - [17] http://www.atomki.hu/TheorPhys/CHSH_activation/
 - [18] A. C. Doherty, P. A. Parrilo, F. M. Spedalieri, Phys. Rev. A **69**, 022308 (2004).
 - [19] B. M. Terhal, A. C. Doherty and D. Schwab, Phys. Rev. Lett. **90**, 157903 (2003).
 - [20] K. F. Pál and T. Vértesi, Phys. Rev. A **82**, 022116 (2010).
 - [21] L. Vandenberghe and S. Boyd, SIAM Review **38**, 49 (1996).
 - [22] D. Collins, N. Gisin, N. Linden, S. Massar, and S. Popescu, Phys. Rev. Lett. **88**, 040404 (2002).
 - [23] Since any two-outcome measurement is a convex combination of von Neumann measurements, w.l.o.g. we can assume the measurement operators to be projectors.